
MAC Equine UltraTM

A nutritional source of Amino Acid Chelated minerals/complex* and vitamins for Horses.



GUARANTEED ANALYSIS

Calcium (max) 12.0 % Phosphorus (min) 6.00 % Magnesium (min) 3.37 % Potassium (min) 1.8 % Colbalt 32 ppm Copper (min) 1000 ppm Magnesse (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin B (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Niacin B-3 (min) 330mg/lb Niacin B-3 (min) 330mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 1200mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb None added	Calcium (min)	9.0 %
Magnesium (min) 3.37 % Potassium (min) 1.8 % Colbalt 32 ppm Copper (min) 1000 ppm Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Jooo I.U./lb 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin B3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Calcium (max)	12.0 %
Potassium (min) 1.8 % Colbalt 32 ppm Copper (min) 1000 ppm Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Methionine (min) 800mg/lb Methionine (min) 130mg/lb Salt None added		
Colbalt 32 ppm Copper (min) 1000 ppm Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin B3 (min) 54,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Magnesium (min)	3.37 %
Colbalt 32 ppm Copper (min) 1000 ppm Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin B3 (min) 54,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Potassium (min)	1.8 %
Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Methionine (min) 800mg/lb Methionine (min) 130mg/lb Salt None added		
Manganese (min) 2600 ppm Selenium yeast (min) 35 ppm Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Methionine (min) 800mg/lb Methionine (min) 130mg/lb Salt None added	Copper (min)	1000 ppm
Zinc (min) 4600 ppm Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 1,70mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 990mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Manganese (min)	2600 ppm
Iodine(min) 10ppm Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Selenium yeast (min)	35 ppm
Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Zinc (min)	4600 ppm
Vitamin A (min) 200,000 I.U./lb Vitamin D3 (min) 54,000 I.U./lb Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg./lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Iodine(min)	10ppm
Vitamin E (min) 3,000 I.U./lb Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Vitamin A (min)	200,000 I.U./lb
Pyridoxine B-6 (min) 320mg/lb Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Vitamin D3 (min)	54,000 I.U./lb
Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Vitamin E (min)	3,000 I.U./lb
Vitamin B-12 (min) 170mg/lb Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Pyridoxine B-6 (min)	320mg/lb
Choline (min) 2600mg/lb Folic acid (min) 53mg/lb Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added		
Niacin B-3 (min) 330mg/lb D-Pantothenic acid (min) 530mg/lb Ascorbic acid C (min) 330mg/lb Biotin (min) 43 mg/lb Riboflavin B-2 (min) 900mg/lb Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added		
D-Pantothenic acid (min) .530mg/lb Ascorbic acid C (min) .330mg/lb Biotin (min) .43 mg/lb Riboflavin B-2 (min) .900mg/lb Thiamine B-1 (min) .1200mg/lb Methionine (min) .800mg/lb Menadione K (min) .130mg/lb Salt .None added	Folic acid (min)	53mg/lb
D-Pantothenic acid (min) .530mg/lb Ascorbic acid C (min) .330mg/lb Biotin (min) .43 mg/lb Riboflavin B-2 (min) .900mg/lb Thiamine B-1 (min) .1200mg/lb Methionine (min) .800mg/lb Menadione K (min) .130mg/lb Salt .None added	Niacin B-3 (min)	330mg/lb
Biotin (min) .43 mg/lb Riboflavin B-2 (min) .900mg/lb Thiamine B-1 (min) .1200mg/lb Methionine (min) .800mg/lb Menadione K (min) .130mg/lb Salt None added		
Biotin (min) .43 mg/lb Riboflavin B-2 (min) .900mg/lb Thiamine B-1 (min) .1200mg/lb Methionine (min) .800mg/lb Menadione K (min) .130mg/lb Salt None added	Ascorbic acid C (min)	330mg/lb
Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added		
Thiamine B-1 (min) 1200mg/lb Methionine (min) 800mg/lb Menadione K (min) 130mg/lb Salt None added	Riboflavin B-2 (min)	900mg/lb
Menadione K (min)130mg/lb SaltNone added		
SaltNone added	Methionine (min)	800mg/lb
SaltNone added	Menadione K (min)	130mg/lb
		None added

Dried Aspergillus Oryzae Fermentation Product (min).....82g/lb

INGREDIENTS

Dicalcium Phosphate, Monocalcium Phosphate, Calcium Amino Acid Chelate*, Yeast Culture, Potassium Amino Acid Complex, Magnesium Amino Acid Chelate*, Zinc Amino Acid Chelate*, Manganese Amino Acid Chelate*, Copper Amino Acid Chelate*, Magnesium Oxide, Calcium Carbonate, Potassium Chloide, Manganous Oxide, Zinc Oxide, Copper sulfate, Dried Aspergillus Oryzae Fermentation Product, Wheat Bran, Cobalt Carbonate, Selenium Yeast, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, di-Methionine, Menadione Sodium Bisulite (source of vitamin K), Riboflavin Supplement, Vitamin B12 Supplement, Biotin, Niacin, Folic Acid, Pyriodoxine HCI, Ascorbic Acid, Thiamine Mononitrate, d-Calcium PantothenateAcid, Chloline Chloride, Ethylenediamine Dihydriodide, Soybean Oil, Natural and Artificial Flavor.

FEEDING DIRECTIONS:

Top Dress on Daily Feed.

Up to 250 lbs. 251-500 lbs. 501-800 lbs. 801-1100 lbs. 1101 lbs. Plus 3/4 OZ. $1\frac{1}{2}$ oz. ½0Z. 1 oz. 2 oz.

*What are minerals?

As simple as it may seem, this question is the first step in examining the role of minerals in nutrition. Webster describes minerals as a solid, crystalline substance (diamond copper, quartz, etc.) not of animal or vegetable origin. Important in this definition is the indication that their origin is not from animal or vegetable source. Minerals for nutritional purpose cannot be synthesized by the body. They must be utilized as natural elements from nature.

*Why are minerals important?

The following table shows various systems in the body which utilize minerals. Research is showing that imbalances or deficiencies in mineral nutrition can affect these systems.

Immune System	Cu, Zn, Fe, Se
Energy Production	Mg, P, Mn
Hormone System	Fe, Mn, Zn, Cu, Mg, K
Vitamin Production	Co
Blood Production	Cu, Fe
Enzyme System	Zn, Cu, Mn, Mg, Fe, Ca, Mo
Skeletal System	Ca, Mg, Zn, Mn, B, P
Reproduction	P, Cu, K, Mn, Zn, Mg

*What is <u>BIOAVAILABILITY</u> and why is it important?

One good definition of **Bioavailability** is the amount of a nutrient ingested that is absorbed and thus available to the body for metabolic use.

Bioavailabilty is important because all nutritional intake must be available to various body systems for growth, maintenance of body tissues, reproduction and other performance factors. No matter how high the nutrient levels or how well formulated the product, if it is not available then money and effort has been wasted.

*What are CHELATED MINERALS?

When minerals such as zinc, manganese, magnesium, copper, iron, calcium, and others become surrounded by and bonded to amino acids, in a stable form, this is referred to as CHELATION. Chelation is a natural means for the body to transport minerals across the intestinal wall as part of digestion.

*Why is it important for the mineral to have a stable bond to the amino acids?

Simply mixing inorganic minerals with amino acids in a liquid or dry mixture does not fall into the category of a true amino acid chelate. This simple ionic and hydrogen bonding of minerals to amino acids does not produce a stable product. Special processing must be preformed to create a stable (covalent) bond which is important for greater bioavailability.

*What makes MAC so affective?

a. Size: Picture in your mind the fuel filter on the car engine. The filter allows fuel to pass through but holds back Large particles from entering the engine. This idea applies to the absorption of minerals from the intestine to the blood stream.

Large particles cannot easily pass through the intestinal wall. Many mineral products on the market have molecular weights too large to be absorbed intact.

Through technology, chelated minerals with molecular weights small enough to pass easily through the intestinal wall. The results is a compound similar to that which the body itself produces by natural chelation.

b. Stability: Stability is maintained throughout the entire ph range encounter during the digestion process. This assures maximum presence of intact mineral chelate for absorption via the special dipeptide-like absorption process.

The body cannot utilize traditional mineral compounds in their natural state. Zinc sulfate, iron sulfate, or any sulfate, oxide or carbonate must be broken apart and restructured to allow it to be transported through the intestinal wall.

A similar situation exists with some reported chelates or complex ed mineral products. Not properly stabilized, they break apart, exposing the raw, ionized mineral.

It is after digestion when other mineral forms have their mineral payload cleaved from their carriers. In this situation, these minerals become charged ions, and their absorbability comes into jeopardy. These charged free minerals are known to block the absorption of one another, or to combine with other dietary factors to form compound that are un absorbable.

c. Neutrality: The process of chelation results in the final mineral compound becoming neutral, i.e., containing no electrical charge. Why is this important? Mineral compounds that have electrical charges can interact with other dietary compounds, such as phytates and other oppositely charged molecules, and form substances that are not absorbable. In addition, mineral compounds that have an electrical charge are reactive, and as such they can deactivate other important nutrient factors, such as: vitamin E, ascorbic acid, various B-vitamins, as well as important medications.

There are many products on the market which are reported to be chelates. Some are only complex mixtures of minerals and proteins not fitting into the definition of a true amino acid chelates. These lose integrity during digestion, becoming unstable and compromising availability. 1

*Why are minerals bound to amino acids to form a chelate?

The body is very efficient at absorbing amino acids. In a priority list of nutritional substance crossing the intestinal wall, after digestion, <u>Dipeptides</u> (two amino acids linked together through a special bond) and single amino acids rank highly. In fact dipeptides now appear to absorbed at a higher rate than a single amino acid. This an apparent by- product of their special active transport mechanism of absorption. It has been seen that dipeptides are removed from the interior of the intestine at a much faster rate than single amino acids. Chelating minerals to amino acids, in a dipeptides-like fashion, allows this mineral form to be smuggled via this special active transport system across the intestinal lining

*Why use MAC Equine Ultra?

Ask yourself, is my horse or horses plagued with various Disease, Disorders or Medical problems; Nervous and Unmanageable; Hoof Problems; Hair & Skin conditions; Breeding Problems; Developmental Growth; Joints; Allergies; etc.

The solution to these problems and many more may be as simple as a change in diet.

If your horse or horses are in good shape. By feeding MAC you will only improve their health and start a preventive medicine program.

MAC is formulated to provide your horse with <u>Ultra Bio-availability</u> of minerals and vitamins by using True Amino Acid Chelates. By using a True Chelated Mineral the horse's body works more efficiently. This will significantly impact their immune system, energy production, hormone system, vitamin production, blood production, enzyme system, skeletal system, reproduction, hoofs, developmental growth, hair and skin conditions, etc.

MAC now has Amaferm®. Amaferm® is an all natural fermentation product produced from a highly selected and specific strain of Aspergillus Oryzae. The result is increased rate of digestion and feed efficiency resulting in increased animal digestive performance.

MAC Representative



MAC Equine Products™

MAĈ Equine Ultra™ MAC Ultra Gel™ MAC Rice Bran Oil MAC Calm MAC Ultra E MAC Ultra Hoof

Arabian Haven Ranch[™] LLC. W261 County Road D Borth, WI 54923 920-987-5693 or 920-252-1057 Arabianhavenranch@gmail.com